

Set Proofs Continued

Set Proofs Continued

Inclusion

Basic principle: $A \subseteq B$ is equivalent to the statement

$x \in A \implies x \in B$. One can prove this both directly and as
 $x \notin B \implies x \notin A$.

Proposition: Let $A = \{4x + 2 : x \in \mathbb{Z}\}$. Let $B = \{2x : x \in \mathbb{Z}\}$.
Then $A \subseteq B$.

$$A = \{4x + 2 : x \in \mathbb{Z}\}$$

$$\rightarrow 2 = 4 \cdot 0 + 2$$

$$-6 \quad 6 \quad \downarrow$$

$$-10 \quad 10$$

$$\vdots \quad 14$$

$$B = \{2x : x \in \mathbb{Z}\}$$

$$\dots -4, -2, 0, 2, 4, 6, 8, \dots$$

Is $A \subseteq B$?

if $x \in A$, then $x \in B$.

$$\text{if } x = 4k + 2 \text{ then } x = 2y$$

$k \in \mathbb{Z} \qquad y \in \mathbb{Z}$

$$\text{Note } x = 2(2k + 1)$$

So $x = 2(y)$ where $y = 2k + 1$
Therefore $x \in B$

More examples

Proposition: For all $k \in \mathbb{Z}$, let $A = \{n \in \mathbb{Z} : n|k\}$ and let $B = \{n \in \mathbb{Z} : n|k^2\}$. Then $A \subseteq B$. (Note: this is problem 3 on page 171.)

$$\text{Let } A = \{n \in \mathbb{Z} : n|k\}$$

$$B = \{n \in \mathbb{Z} : n|k^2\}$$

k is fixed at the beginning

$$\text{For all } k \in \mathbb{Z}, n|k \Rightarrow n|k^2$$

Since $n|k$, $k = nd$.

$$k^2 = n^2 d^2 = n(nd^2)$$

$$= ny \text{ where } y = nd^2.$$

so $n|k^2$.

this shows $A \subseteq B$.

More examples

Proposition: Suppose A , B , and C are sets. If $B \subseteq C$, then $A \times B \subseteq A \times C$. (This is problem 7 on page 171.)

Suppose A, B, C are any sets.

if $B \subseteq C$, then $A \times B \subseteq A \times C$.

if $\underline{[x \in B \Rightarrow x \in C]}$ then $\underline{[x \in A \times B \Rightarrow x \in A \times C]}$

Proof: Assume $x \in B \Rightarrow x \in C$. ($B \subseteq C$).

and we assume $x \in A \times B$.

$x = (a, b)$ where $a \in A$ and $b \in B$.

We must show $(a, b) \in A \times C$.

in other words, ~~(a, b)~~ we must show

that $(a, b) = (a', c')$ where $a' \in A$
 $c' \in C$

since $b \in B$, $b \in C$ by hypothesis $B \subseteq C$. Therefore $c' = b$
 $(a, b) \in A \times C$, $a' = a$

One more

Proposition: Let A and B be sets. Prove that $A \subseteq B$ if and only if $A - B = \emptyset$. (This is problem 21 on page 171.)

$$A - B = \{x \in A : x \notin B\}$$

$$x \in A - B \iff x \in A \text{ and } x \notin B$$

$A - B = \emptyset \iff$ there does not exist x such that $x \in A$ and $x \notin B$.

NOT [there exists x such that $x \in A$ and $x \notin B$]



For all x , $x \notin A$ or $x \in B$
NOT ($x \in A$ and $x \notin B$)

For all x , $x \notin A$ or $x \in B$.

$A \subseteq B \iff$ for all x , $x \in A \Rightarrow x \in B$.

For all x ($x \in A \Rightarrow x \in B$) \iff for all x ($x \in B$ or NOT $x \in A$)

P	Q	$P \Rightarrow Q$	Q or NOT P
T	T	T	T
T	F	F	F
F	T	T	T
F	F	T	T

$A \subseteq B$ $A - B = \emptyset$


 For all x , $x \in B$ or $x \notin A$.